

# Significance of Perylene for Source Allocation of Terrigenous Organic Matter in Aquatic Sediments

Ulrich M. Hanke,<sup>\*,†,Ⓛ</sup> Ana L. Lima-Braun,<sup>†,||</sup> Timothy I. Eglinton,<sup>†,‡</sup> Jeffrey P. Donnelly,<sup>§</sup> Valier Galy,<sup>†</sup> Pascale Poussart,<sup>†,⊥</sup> Konrad Hughen,<sup>†</sup> Ann P. McNichol,<sup>§</sup> Li Xu,<sup>§</sup> and Christopher M. Reddy<sup>†,Ⓛ</sup>


<sup>†</sup>Department of Marine Chemistry and Geochemistry, Woods Hole Oceanographic Institution, 266 Woods Hole Road, Woods Hole, Massachusetts 02543, United States

<sup>‡</sup>Geological Institute, ETH Zürich, Sonneggstrasse 5, 8092 Zurich, Switzerland

<sup>§</sup>Department of Geology and Geophysics, Woods Hole Oceanographic Institution, 266 Woods Hole Road, Woods Hole, Massachusetts 02543, United States

## Supporting Information

**ABSTRACT:** Perylene is a frequently abundant, and sometimes the only polycyclic aromatic hydrocarbon (PAH) in aquatic sediments, but its origin has been subject of a longstanding debate in geochemical research and pollutant forensics because its historical record differs markedly from typical anthropogenic PAHs. Here we investigate whether perylene serves as a source-specific molecular marker of fungal activity in forest soils. We use a well-characterized sedimentary record (1735–1999) from the anoxic-bottom waters of the Pettaquamscutt River basin, RI to examine mass accumulation rates and isotope records of perylene, and compare them with total organic carbon and the anthropogenic PAH fluoranthene. We support our arguments with radiocarbon (<sup>14</sup>C) data of higher plant leaf-wax *n*-alkanoic acids. Isotope-mass balance-calculations of perylene and *n*-alkanoic acids indicate that ~40% of sedimentary organic matter is of terrestrial origin. Further, both terrestrial markers are pre-aged on millennial time-scales prior to burial in sediments and are insensitive to elevated <sup>14</sup>C concentrations following nuclear weapons testing in the mid-20<sup>th</sup> Century. Instead, changes coincide with enhanced erosional flux during urban sprawl. These findings suggest that perylene is definitely a product of soil-derived fungi, and a powerful chemical tracer to study the spatial and temporal connectivity between terrestrial and aquatic environments.



## INTRODUCTION

Perylene is found in marine<sup>1–5</sup> and lacustrine sediments,<sup>6–8</sup> soils,<sup>9,10</sup> and also petroleum<sup>11,12</sup> and fuel emissions<sup>13,14</sup> often associated with other distinctive anthropogenic combustion-derived polycyclic aromatic hydrocarbons (PAHs). However, most studies report that sediment records of perylene differ greatly from those of anthropogenic PAHs.<sup>15–19</sup> The latter are typically most abundant in sediments post-dating the Industrial Revolution, particularly those deposited during the latter half of the 20<sup>th</sup> Century, whereas perylene abundances are lowest near the sediment–water interface and tend to increase with depth, particularly in anoxic sedimentary settings.<sup>18,20</sup> This depth-related increase in concentration implies abiotic or biologically mediated in situ production from precursor natural product(s)<sup>2,21</sup> by either a first- or second-order reaction under anaerobic conditions.<sup>20</sup>

The origin of perylene has remained unclear for decades, during which has previously been argued for terrestrial<sup>1,2,22,23</sup> and diagenetic,<sup>6,16,24</sup> as well as petrogenic<sup>12,25</sup> and pyrogenic<sup>13,14</sup> sources. It was only recently that Itoh and collaborators<sup>23</sup>

determined that the fungal species *Cenococcum geophilum* produces 4,9-dihydroxyperylene-3,10-quinone, confirming a longstanding hypothesis<sup>1,22</sup> that there is a naturally produced precursor. This ectomycorrhizal fungus appears almost ubiquitous in boreal, temperate, and subtropical regions, and is present in the rhizosphere of woody-plant roots and more generally in forest soils.<sup>26</sup> A fungal origin can explain the widespread abundance of perylene in the environment even over geological time-scales,<sup>27</sup> since mycorrhizal associations with vascular plants evolved about 400 million years ago.<sup>28</sup>

Besides *C. geophilum*, other mycorrhiza also produce similar perylene precursor compounds<sup>29,30</sup> as toxins involved in pathogenesis of their host plant.<sup>31</sup> Prior stable carbon isotopic ( $\delta^{13}\text{C}$ ) measurements of perylene found values similar to those of terrestrial sources ( $\text{C}_3$ -vegetation:  $\delta^{13}\text{C} \approx -27 \text{‰}$ ),

Received: April 17, 2019

Revised: June 11, 2019

Accepted: June 19, 2019

Published: June 19, 2019

supporting a wood-degrading origin from the rhizosphere.<sup>32</sup> This is further supported by dual-isotope analyses of perylene ( $\delta^{13}\text{C}$  and  $\delta\text{D}$ ) in sediments showing similar  $\delta\text{D}$  values as methoxy groups in lignin while  $\delta^{13}\text{C}$  values are consistent with the expected fractionation range in saprophytic fungi.<sup>22</sup> In addition, natural abundance radiocarbon ( $^{14}\text{C}$ ) analyses allows a comparison of the  $^{14}\text{C}$  age of perylene with that of other organic matter constituents,<sup>25</sup> including total organic carbon (TOC) and specific markers of terrigenous organic matter co-deposited in aquatic sediments. Such analyses shed light on the origin and provide further evidence on the origin of perylene and its use in environmental forensics.

Our motivation is to reconcile existing hypotheses on the source of perylene and assess the potential of this marker compound as a biogeochemical tracer to follow the trajectories of terrigenous organic matter mobilization and transport within watersheds. Specifically, we investigate whether it serves as a molecular marker for rhizosphere carbon from catchment soils and thus facilitates source approximation of terrigenous organic matter in sediments. Moreover, while researchers currently tend to exclude perylene from forensic investigations involving the apportionment of PAHs due to its incongruent behavior, perylene may have mutagenic effects on organisms.<sup>33,34</sup> Thus, an improved understanding of the provenance and dynamics of perylene may be pertinent to mapping inventories of natural and anthropogenic pollutants, delineating transport pathways, and reconstructing historical land-use.

In this study, we construct historical records of perylene and TOC abundances and isotope compositions from the anoxic sediments of the suburban Pettaquamscutt River basin, RI. We chose these sediments because they provide an exceptionally detailed chronological record and a wealth of background information on the catchment area, including human influence on local to regional scales.<sup>15,35–38</sup> In addition, this sedimentary record extends over 260 years, from the preindustrial (~1735) until 1999, including an interval characterized by frequent above-ground nuclear-bomb testing (resulting in elevated atmospheric  $^{14}\text{C}$  concentrations peaking in 1963 when the testing ceased), thus offering the opportunity to study the rate at which specific organic carbon species incorporate atmospheric  $\text{CO}_2$  and are sequestered in aquatic sediments. These characteristics of the study site, when coupled with down-core  $\delta^{13}\text{C}$  and  $^{14}\text{C}$  records, provide a novel perspective on perylene biogeochemistry, yielding constraints on its source and on the burial efficiency of terrigenous organic matter in aquatic sediments. We compare mass accumulation rates (MAR) of perylene and TOC to the combustion-derived PAH marker fluoranthene. While several nonalkylated PAHs could be used to trace past-combustion practices, we choose fluoranthene because its quantitative down-core profile resembles the temporal evolution of the sum of 15 parent PAHs in this watershed.<sup>15</sup> We further support our interpretation with down-core data of particle size and leaf wax ( $\text{C}_{30-32}$ ; *n*-alkanoic acids)  $^{14}\text{C}$  variations. We then carefully assess whether the rhizosphere of wooded land serves as the direct source for perylene in the environment and discuss the process of pre-aging of terrestrial organic matter prior to delivery into the aquatic environment in the context of parallel sedimentary records.

## ■ EXPERIMENTAL SECTION

A series of seven freeze cores were collected in the depocenter of the lower basin of the Pettaquamscutt River basin, RI

(41.503100; -71.450500) in 1999. The catchment area covers 35 km<sup>2</sup> of forest, wetlands and open water of which today about 30% is residential land.<sup>37</sup> Ocean water flooded the basin about 1700 ± 300 years ago leading to a stratified water column and sustained anoxic conditions in bottom waters and underlying sediments.<sup>36</sup> A detailed description on the sediment chronology and sample processing is presented elsewhere.<sup>15,39</sup> Sediment chronology was obtained from varve counting of X-ray radiographs as well as from  $^{137}\text{Cs}$  and  $^{210}\text{Pb}$  profiles using the model of constant rate of supply that yielded a sequence of about 260 years (1735–1999 AD) and an average sedimentation rate of  $0.44 \pm 0.10 \text{ cm yr}^{-1}$ .<sup>39</sup>


Sections from the cores were combined after aligning x-radiograph images including hurricane layers in 1938 and 1954 to a reference chronology of varve counting,  $^{210}\text{Pb}$ , and  $^{137}\text{Cs}$ . This was necessary to obtain enough sediment for trace molecular isotopic-analyses.<sup>35</sup> Hence, we pooled the samples in eight horizons for all isotopic analyses: H1 (1999–1982), H2 (1981–1962), H3 (1960–1931), H4 (1929–1898), H5 (1896–1873), H6 (1871–1842), H7 (1840–1768), and H8 (1764–1735), respectively, except *n*-alkanoic acids that were analyzed in four individual samples:  $1885 \pm 5$ ,  $1947 \pm 33$ ,  $1972 \pm 2$ , and  $1991 \pm 1$ . Quantitative measurements for the calculation of the mass accumulation rates as well as grain size analyses were performed on individual samples as well.

**Grain Size Analyses.** One freeze-core slab was subsectioned using a scalpel blade, transferred into 50 mL round-bottom flasks and treated with a 30% hydrogen peroxide solution at a ratio of 40 mL per 1 g sediment, and heated in a water bath to 70 °C as a means to support mineralization of the organic matter. Subsequently, samples were freeze-dried, and a subsample of about 11 mg was suspended in water and measured on a Beckmann Coulter LS13 320 laser diffraction particle analyzer (Indianapolis, IN) in triplicate.

**Total Organic Carbon and Nitrogen.** A Fisons 1108 elemental analyzer was used to measure the TOC content of the samples. To remove the inorganic carbon fraction, about 2 mg of dry sample was weighed into a silver capsule and acidified with 20  $\mu\text{L}$  of 2 N HCl. TOC content was calculated in relation to the whole sediment dry weight while organic carbon/organic nitrogen TOC/TN ratios were calculated on an atomic basis. Samples were run in triplicate and all reported weight percentages represent the mean ± one standard deviation with an instrumental blank of 0.004 mg for C and smaller than 0.005 mg for N.

**Isotope Ratio Monitoring Mass Spectrometry.** The stable carbon isotopic composition of bulk sample TOC was determined in triplicate by automated online solid combustion interfaced to a Finnigan Delta Plus isotope ratio mass spectrometer. Isotope ratios were calculated relative to  $\text{CO}_2$  reference gas pulses, with standard deviations for replicate measurements were always better than 0.6‰ and usually within 0.3‰.

**Extraction, Purification, and Isotope Analyses of Perylene, Fluoranthene, and *n*-Alkanoic Acids.** Dry sediment samples (0.5–1.5 g) were extracted by pressurized fluid extraction (Dionex ASE 200) using a mixture of acetone and *n*-hexane (1:1) at 1000 psi at 100 °C. Molecular identification and quantification was achieved using an Agilent 6890 Plus GC System interfaced to a mass selective detector operating at 70 eV in SIM mode using a DB-XLB capillary column (60 m × 0.25 mm × 0.25  $\mu\text{m}$ ).<sup>15</sup>



**Figure 1.** Mass accumulation rates for total organic carbon (TOC), perylene, and fluoranthene as well as carbon to nitrogen (TOC/TN) ratios and down-core shifts in coarse particle abundance ( $D_{90}$ ) with shaded area A representing the maximum PAH flux in this area, B the period of amplified erosion due to infrastructure development, and C the preindustrial era prior to 1840s.

Compound-specific radiocarbon analyses (CSRA) of perylene and fluoranthene were performed on eight pooled horizons by first using high-pressure liquid-chromatography (HPLC) to separate pure perylene (98% purity or greater) from the sample extracts. The HPLC procedure isolated PAHs into 2-ring and the combined 3 + 4-ring and 5 + 6-ring PAHs. The resulting 16 HPLC fractions (8 horizons  $\times$  2 ring classes) were subjected to two-dimensional preparative capillary gas chromatography for isolation and purification of individual PAHs via HP 7683 autinjector and a multicolumn switching system (Gerstel MCS 2) connected to a HP 6890 series gas chromatograph, and Gerstel preparative fraction collector (PFC; further details in the [Supporting Information \(SI\)](#)). Purified samples were transferred to precombusted quartz tubes (7 mm i.d.  $\times$  20 cm), dried under nitrogen before adding copper oxide (50 mg). Each tube was then evacuated on a vacuum line while samples were kept at  $-90$  °C to prevent sublimation, sealed, and combusted at 850 °C for 5 h. About 95% of the purified carbon dioxide was reduced to graphite, pressed and analyzed for  $^{14}\text{C}$  at NOSAMS, Woods Hole, MA, and the remaining 5% was used for  $\delta^{13}\text{C}$  measurements.


The  $\delta^{13}\text{C}$  values of perylene were also determined in triplicate on a Finnigan Delta Plus isotope ratio mass spectrometer with attached Finnigan GC combustion III interface and Hewlett-Packard 6890 GC (irm-GC/MS). Compounds were separated on a CP-Sil 5CB capillary column (50 m  $\times$  0.25 mm  $\times$  0.25  $\mu\text{m}$ ) and isotope ratios for PAH peaks were calculated relative to  $\text{CO}_2$  reference gas pulses. The standard deviation for replicate measurements of perylene was better than 0.6 ‰ and mostly around 0.2 ‰.

The isotopic composition of leaf wax *n*-alkanoic acids ( $\text{C}_{30-32}$ ) were determined for four individual sediment samples deposited in  $1885 \pm 5$ ,  $1947 \pm 33$ ,  $1972 \pm 2$ , and  $1991 \pm 1$  following the analytical procedure described elsewhere.<sup>40</sup> In brief, *n*-alkanoic acids were extracted, isolated, and purified using a preparative Hewlett-Packard 5890 series II capillary gas

chromatograph.<sup>41</sup> Following the chromatographic purification of individual compounds, samples were graphitized and measured at NOSAMS, Woods Hole, MA.

**Evaluation of Isotope Data.** Isotope mass-balance calculations provide quantitative estimates for source apportionment of organic matter. To constrain the sources of TOC and perylene in Pettaquamscutt River sediments, we calculated the relative contribution of possible organic carbon sources in the sediments using the measured  $^{14}\text{C}$  and  $\delta^{13}\text{C}$  values for TOC (Figure 2). We report the  $^{14}\text{C}$  data as  $F^{14}\text{C}$  which is the fraction modern independent from the year of measurement.<sup>42</sup> We assumed a simple mixing model to retrieve an average value for the time lag of perylene and leaf wax *n*-alkanoic acids; the details of which are published elsewhere.<sup>35</sup> In brief, these source-specific molecular markers facilitate approximations of the contributions of terrigenous organic matter in aquatic sediments (further details on the concept are included in the [SI](#)).

To account for the potential variability under increasing human pressure on the coastal environment, we determined average values of our terrestrial end-members for two time periods: the preindustrial (1735–1840) and the post-“bomb” (1960–1999) period (see [SI](#) for more detail). Regarding  $^{14}\text{C}$  data evaluation, the post-1960s period also is a benchmark to test for the resilience and short-term dynamics of catchment processes. In contrast, the aquatic end-member is only affected by the marine reservoir effect, which is about  $400 \pm 40$   $^{14}\text{C}$  years lower than the atmospheric  $^{14}\text{C}$  concentration at the time of burial.<sup>43</sup> Furthermore, we neglect a potential impact of petrogenic carbon<sup>44</sup> in the Pettaquamscutt River because of the absence of postglacial erosion and due to the absence of carbonaceous metasedimentary rocks in the basin.<sup>45</sup> We constrain our model by  $\delta^{13}\text{C} = -21.0$  ‰ for aquatic production<sup>46</sup> and use local records of perylene, pyrogenic carbon<sup>35</sup> and *n*-alkanoic acids ( $\text{C}_{30-32}$ ) to define terrestrial end-members. We further obtained the contemporary atmospheric



**Figure 2.** High-resolution  $\delta^{13}\text{C}$  record for TOC and pooled sample results for perylene (left); molecular  $^{14}\text{C}$  data for the combustion-derived fluoranthene (middle) and results on perylene, leaf wax *n*-alkanoic acids as well as TOC (right).

$^{14}\text{C}$  concentrations from the extended Intcal13 reference chronology.<sup>43,47</sup>

## RESULTS AND DISCUSSION

The exceptionally well-constrained chronology of the Pettaquamscutt River sediments along with detailed reconstructions of the provenance of combustion-derived PAHs<sup>15,35</sup> and  $\text{Hg}^{38}$  provide key constraints that allow us to determine whether perylene signatures are consistent with its production in the rhizosphere of catchment soils.<sup>23,26</sup> To characterize the export of terrigenous organic matter, including the putative precursors of perylene, and its subsequent burial in aquatic sediments, we report MAR for TOC, perylene, and fluoranthene. Isotopic mass balance calculations for  $\delta^{13}\text{C}$  and  $^{14}\text{C}$  records are employed to further constrain the main source of perylene precursors. With reference to recent research and our results, we then provide a synthesis aiming to merge the existing concepts, reconciling the origin of perylene (i.e., terrestrial, diagenetic or petro-/pyrogenic).

**Down-Core Mass Accumulation Rates of TOC and Perylene.** The burial rate of TOC in the sediments of the Pettaquamscutt River ranges from 38 to 61  $\text{g m}^{-2} \text{yr}^{-1}$  (average =  $49 \pm 6 \text{ g m}^{-2} \text{yr}^{-1}$ ,  $n = 71$ ). The lowest TOC burial flux is observed in ca. 1938 (Figure 1) and is related to a 16.7% shift in burial flux from  $53.0 \pm 43 \text{ g m}^{-2} \text{yr}^{-1}$  prior to ca. 1927 to  $44.0 \pm 3.7 \text{ g m}^{-2} \text{yr}^{-1}$  for sediments deposited thereafter. The timing of this change coincides with the construction of the Lacey Bridge on the ocean side of the lower basin in 1934<sup>45</sup> (Figure 1B) that likely caused a restriction in seawater intrusion, affecting the influx of both marine as well as the terrestrial matter. We argue that this exemplifies the accelerated development of the catchment area beginning early in the last century and the related infrastructure projects that changed the natural flow paths of materials within the watershed.

The  $N_{\text{org}}$  down-core record reveals only small variations, here illustrated as  $C_{\text{org}}/N_{\text{org}}$  profiles ( $n = 71$ ; Figure 1) where three trends are delineated: decreasing C/N from 1730s (12.6) to 1840s (9.4), a  $\sim 50$ -year reversal toward higher C/N ratios (13.2 in 1900), and a century-long C/N secular decrease in the upper core (10.0 in 1999). This rather narrow range in C/N ratios suggests that organic matter is predominantly yet not exclusively

composed of labile material with C/N values similar to aquatic organic matter (C/N = 4 to 10).<sup>48</sup> However, we refrain from further speculations on this bulk-level information due to the absence of a robust reference for C/N values of terrigenous organic matter from this area.

MARs for perylene (and proto-perylene), which corresponds to the reduced perylenequinone, vary from 29 to 3800  $\mu\text{g m}^{-2} \text{yr}^{-1}$ , while its down-core profile differs markedly from the combustion-marker fluoranthene (Figure 1). The values were low before 1938, reached a maximum between ca. 1938 and 1973 ( $1580 \mu\text{g m}^{-2} \text{yr}^{-1}$ ), before diminishing again in most recent decades ( $\sim 100 \mu\text{g m}^{-2} \text{yr}^{-1}$ ) (Figure 1). This contrasting accumulation behavior implies a different origin for perylene relative to combustion-derived PAHs, here illustrated by fluoranthene (Figure 1). This nonalkylated PAH appears lower in abundance varying from 1 to 1280  $\mu\text{g m}^{-2} \text{yr}^{-1}$  and also exhibits significant variability,<sup>35</sup> however the temporal evolution of MARs of this pyrogenic PAH differs sharply from that of perylene. Specifically, fluoranthene's MAR was low until the mid-19th century (Figure 1C), increased steadily to a maximum in the late-1950s, then declined in the post-1970s due to cleaner burning fuels and as consequence of environmental regulations.<sup>15,37</sup> Such down-core profiles of combustion-derived PAHs are also observed in other sedimentary records, including the upper basin of the Pettaquamscutt River.<sup>16,19,49</sup>

The observed decline in perylene (and proto-perylene) MAR between 1927 and 1938 coincides with the abrupt decrease in TOC accumulation rate (Figure 1B). We attribute the latter to a dilution in organic carbon content resulting from a higher proportion of clastic material due to erosion of soil mineral horizons (Figure 1). Perylene is a trace constituent of TOC, never accounting for more than 0.1% of TOC, and consequently perylene/TOC ratios shift by 500% relative to before or after this time interval. These observations thus support the link between a decrease in perylene's MAR resulting from an increase in erosional flux (Figure 1B).

A striking feature of the perylene profile is that fluxes remain nearly constant ( $665 \pm 180 \mu\text{g m}^{-2} \text{yr}^{-1}$ ;  $n = 26$ ) prior to 1865, indicating either steady-state erosional conditions or quantitative conversion of precursor material (Figure 1).<sup>23</sup> Due to the large variations in perylene burial rates in our sedimentary

record we refrain from attempting to derive kinetic parameters for conversion of perylenequinone (proto-perylene) precursor to perylene. However, Slater and collaborators<sup>50</sup> compared two perylene profiles from Lake Siskiwit collected in 1983 and 2005. They found a reaction rate constant ( $0.048 \text{ yr}^{-1}$ ), ca. 75% higher than previously reported,<sup>20</sup> whereas another study reported much lower rates.<sup>18</sup> All of the above studies suggest that the conversion of perylenequinone to perylene follows first-order kinetics. Nevertheless, the strong ( $\sim 97\%$ ) decrease in perylene abundance in Pettaquamscutt River sediments from 1973 to 1999 likely reflects incomplete conversion of precursor compounds (Figure 1).

**Isotope Mass Balance Calculations.** The  $\delta^{13}\text{C}$  TOC profile (Figure 2) appears relatively invariant throughout the core ( $-25.4$  to  $-23.0\%$ , average  $-24.1 \pm 0.5\%$ ,  $n = 36$ ), indicating the absence of large variations in organic matter composition. Although the TOC MAR decreased in the 1930s,  $\delta^{13}\text{C}$  values do not suggest this was accompanied by a shift in the nature of organic matter supplied to Pettaquamscutt River basin. Instead, the observed changes in burial flux in this catchment are attributed to a shift in land use stemming from an increase in the local human population and the accompanied development of infrastructure. The  $\delta^{13}\text{C}$  values of perylene ( $-28.9$  to  $-28.3\%$ ; average  $-28.6 \pm 0.6\%$ ; Figure 2) were similarly invariant, while significantly lower than TOC and similar to that reported for terrestrial  $\text{C}_3$  plants ( $-29$  to  $-25$ , average  $-28\%$ ).<sup>51</sup>

In contrast, the  $^{14}\text{C}$  record of TOC for pooled samples ( $n = 8$ ; Figure 2) reveals only small variations in preindustrial time (1735–1840) but then tends to shift toward more  $^{14}\text{C}$  depleted values between 1839 and 1958. This change is thought to integrate the increasing use of fossil fuels slightly diluting the natural atmospheric  $^{14}\text{C}$  concentration (the so-called “Suess effect”) as well as ongoing infrastructural development and constructions within the catchment resulting in the mobilization of deeper mineral soils. Thereafter, we observed a drastic ( $\sim 15\%$ ) increase in  $\text{F}^{14}\text{C}$  TOC values reflecting the shift in atmospheric  $^{14}\text{C}$  derived from thermonuclear weapons testing in the post-1960s. The uppermost sediment layer signals declining  $^{14}\text{C}$  concentrations similar to the atmosphere.<sup>47</sup>

Compound-specific radiocarbon analysis reveals that the  $^{14}\text{C}$  profile of perylene follows that of TOC from the preindustrial period until  $\sim 1950$ . Thereafter,  $^{14}\text{C}$  of TOC increases due to the incorporation of atmospheric bomb  $^{14}\text{C}$ , while perylene tends to more  $^{14}\text{C}$ -depleted values. This could reflect a pyrogenic (fossil fuel) contribution to perylene, mitigating the impact of “bomb” carbon in the Pettaquamscutt River sediments. However, CSRA of leaf wax *n*-alkanoic acids ( $\text{C}_{30-32}$ ) from the same sediments reveals a concomitant  $^{14}\text{C}$  trend with perylene (Figure 2). We infer, therefore, that leaf waxes and perylene share a common source and mode of export, implying that local erosional fluxes and mobilization of soil organic matter are responsible for the decoupled TOC and perylene post-1950s signals rather than global-scale perturbations such as “Suess effect” or above-ground nuclear weapons testing.

In the mainly wooded catchment area of Pettaquamscutt River, organic molecules are produced during photosynthetic activity, translocated and metabolized in plants, as well as released into the rhizosphere. In soil, plants and animals, residues can be assimilated, stabilized and metabolized until they eventually become respired to carbon dioxide,<sup>52</sup> or they can be eroded and redistributed within the watershed.<sup>53</sup> Source-specific molecular markers allow delineation of specific processes associated with organic matter cycling, while molecular  $^{14}\text{C}$

data provide additional information on the time scales of organic matter storage and transport from its source to the “ultimate” site of burial. This “residence time” has been determined for several markers of terrestrial primary productivity,<sup>54–56</sup> as well as charred plant biomass,<sup>35,57</sup> and range from centuries to several millennia. The underlying processes resulting in organic matter “pre-aging” prior to burial in aquatic depocenters can complicate source apportionment because the  $^{14}\text{C}$  concentration of various organic matter components differs from that of the corresponding atmospheric  $^{14}\text{C}$  reference value.

We determined an average  $^{14}\text{C}$  residence time for perylene of  $1300 \pm 300$   $^{14}\text{C}$  years for samples pre-dating the industrialization (1735–1840,  $n = 3$ ) and  $2000 \pm 500$   $^{14}\text{C}$  years for the post-“bomb” era (1960–1999;  $n = 3$ ). These values agree well with leaf wax *n*-alkanoic acids ( $\text{C}_{30-32}$ ) that showed a  $2700 \pm 700$   $^{14}\text{C}$  year residence time for the post-“bomb” era (Figure 2), and a record of biomass-derived pyrogenic carbon from the same core revealed an average age of  $1460 \pm 490$   $^{14}\text{C}$  years for the preindustrial era.<sup>35</sup> The apparent average age or residence time over the preindustrial and the post-“bomb” era for perylene and higher plant *n*-alkanoic acids of ca. 2000  $^{14}\text{C}$  years is slightly higher than it has been reported for (micro) charcoal in the same catchment<sup>35</sup> as well as for leaf wax *n*-alkanoic acids in the Bengal fan<sup>55</sup> yet similar to that of Cariaco basin.<sup>56</sup> However, in the post-“bomb” era we observed a decoupling of  $^{14}\text{C}$  concentrations among molecular markers and TOC. Specifically, the increase in atmospheric  $^{14}\text{C}$  concentrations post-1960s is not apparent in perylene or *n*-alkanoic acids, suggesting that soil carbon storage on millennial time scales may conceal or delay the legacy.

Isotope mass balance calculations based on TOC  $^{13}\text{C}$  and  $^{14}\text{C}$  data yield similar values for the proportion of organic matter of terrestrial origin ( $\delta^{13}\text{C}$ :  $-43 \pm 10\%$  and  $^{14}\text{C}$ :  $-37 \pm 10\%$ ), with the remaining carbon derived from aquatic productivity. These results for the Pettaquamscutt River basin are slightly higher than an estimated global average of one-third of sedimentary organic is of terrestrial origin.<sup>58</sup> The similar estimates obtained from dual-carbon isotope evaluations based on perylene reinforce its value in constraining the terrestrial end-member in source apportionment.

**New Constraints on the Source of Perylene.** The origin and widespread distribution of perylene in the environment has been a subject of scientific debate for decades, although its conversion from the precursor molecule, 4,9-dihydroxyperylene-3,10-quinone, has long been suspected.<sup>1,2,23,29</sup> In particular, it has remained unclear whether perylene is formed during *in situ* diagenesis<sup>6,16</sup> or is of petro-/pyrogenic origin.<sup>11,14,25,59</sup> The former hypothesis was stimulated by the presence of perylene in Antarctic marine sediments<sup>60</sup> and by large similarities between perylene and total organic matter down-core profiles, fueling the notion of *in situ* synthesis from TOC.<sup>16</sup> Along these lines, Gschwend and collaborators<sup>20</sup> calculated kinetic parameters necessary to yield perylene from biogenic precursor concentrations that are consistent with results from a recent study comparing down-core profiles 20 years apart in Lake Siskiwit,<sup>50</sup> and both are in line with the observed reduction of perylenequinones.<sup>23</sup> The second alternative hypothesis is based on the abundance of perylene in ancient sediments,<sup>22,61</sup> fossil fuels<sup>11,12</sup> or combustion emissions.<sup>13,14,59</sup> Regarding the latter sources, perylene concentrations remain below 1.4% of the total PAH and thus are negligible for the overall budget. Its presence in these matrices, however, raises the question of whether perylene is exclusively of fungal origin. A possible explanation for the

occurrence of perylene in fossil fuels lies in the evolutionary development of fungi. The symbiotic interaction between fungi and vascular plants evolved about 400 Ma ago, and ectomycorrhiza have existed for at least 56 Ma,<sup>62</sup> while the divergence and evolution of the mycorrhiza gene pool remains largely unexplored.<sup>28</sup> However, Blumer proposed the conversion of pigments to hydrocarbons through a geochemically irreversible deoxygenation of functional groups and hydrogenation under sustained reducing conditions.<sup>63</sup> Even though this was exemplified for fossil crinoids in a pigment-rich Triassic oil shale, these reactions likely also apply to the precursors of biogenic PAH. We deduce, therefore, that fungal-derived perylenequinone is deoxygenated during sediment diagenesis and, at least partly, survives coalification. This could explain the occurrence of perylene in fossil deposits<sup>32</sup> and could explain its abundance in combustion residues. One evolutionary rationale for the occurrence of perylene in sclerotia, the resting structures of *C. geophyllum*,<sup>64</sup> is the extreme mutagenic activity of the unsaturated nucleus of perylenequinones on Gram-negative bacteria,<sup>33</sup> protecting the reproductive function of fungal spores. Overall, the preponderance of evidence,<sup>17,22,32</sup> reinforced by our new data, indicates that perylene is a remnant of wood degradation or fungal activity in the rhizosphere, a hotspot of biological activity on land, and source of organic matter to aquatic systems.

**Parallels between Perylene and TOC.** Sedimentary organic matter is a highly complex mixture composed of aquatic and terrigenous organic carbon, and this complexity complicates source apportionment calculations, as well as determination of carbon fluxes and thus the carbon burial efficiencies.<sup>65,66</sup> Measurement of the abundance and isotopic composition of source-specific molecular markers can provide constraints on terrigenous organic matter in sediments.<sup>65</sup> In addition to source constraints from  $\delta^{13}\text{C}$  and  $^{14}\text{C}$  signatures, the latter also yields information on whether organic matter is of modern or fossil origin,<sup>44</sup> as well as on time scales of terrigenous organic matter between formation on land and burial in aquatic depocenters (i.e., “average residence times”).<sup>35</sup> The process of “pre-aging” implies a time lag between biosynthesis and export of organic matter from land to depocenter, with natural as well as anthropogenic molecules being retained in catchment soils from annual to millennial time-scales.<sup>35,55,56</sup> Here, we find that this pre-aging process occurs on similar time-scales for leaf wax fatty acids (*n*-alkanoic acids), biomass-derived pyrogenic carbon and the rhizosphere tracer perylene—all together terrestrial markers with different modes of formation, chemical structures and functionalities. This agreement between these different source-specific markers suggests that retention within soils regulates the export of most terrigenous organic matter and associated carbon-based pollutants that enter the soil column.

TOC and perylene MARs covary in the preindustrial era until about 1850 despite some variability in the TOC flux (Figure 1). We attribute the latter to the increasing pressure by extensive farming, as well as potential changes in aquatic productivity. The dual carbon isotope records for TOC and perylene (Figure 2) support the absence of drastic compositional shifts throughout the record, despite the influence of the “Suess effect” and nuclear weapons testing on the latter part of the TOC  $^{14}\text{C}$  record. Notably, perylene shows large variation in MAR after 1850, which we suspect is due to the conversion of the catchment from a rural into a suburban landscape that was accompanied by an enhanced erosional flux of carbon from bare soils.

Similarities between TOC and perylene records have been observed previously,<sup>6,16,22,61</sup> yet in these cases it was suspected to reflect direct in situ microbial production of perylene<sup>16</sup> rather than the reduction and conversion of a precursor into perylene. This depletion of the precursor pool may continue over geological time scales,<sup>63</sup> thus limiting comprehensive quantitative assessment of perylene in recent sediments.<sup>50</sup> In contrast, the isotopic composition of perylene is expected to be insensitive to conversion efficiency, and thus can serve as a robust tracer of soil-derived terrestrial organic matter.

## ■ ASSOCIATED CONTENT

### 📄 Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.est.9b02344.

Details on the preparative chromatographic isolation of PAHs for compound-specific isotope analysis, the concept of two modes of terrigenous organic matter export from land to oceans, the impact of the nuclear weapon testing on the atmospheric and marine  $^{14}\text{C}$  partitioning over time and results for grain size analyses, respectively (PDF)

## ■ AUTHOR INFORMATION

### Corresponding Author

\*E-mail: uhanke@whoi.edu.

### ORCID

Ulrich M. Hanke: 0000-0002-4270-225X

Christopher M. Reddy: 0000-0002-7814-2071

### Present Addresses

<sup>||</sup>(A.L.L.B.) ExxonMobil Exploration Company, Spring, Texas, United States.

<sup>⊥</sup>(P.P.) Princeton University, 36 University Place, Princeton, New Jersey 08544, United States.

### Notes

The authors declare no competing financial interest.

## ■ ACKNOWLEDGMENTS

We thank John King, Sean Sylva, Brad Hubeny, Peter Sauer, and Jim Broda for their help in sampling; Carl Johnson and Daniel Montluçon for their incessant help with analyses; as well as Mark Yunker for critical discussion on the perils of perylene. Professor Phil Meyers and two anonymous reviewers provided comments that improved the quality of the manuscript. U.M.H. acknowledges the Swiss National Science Foundation for his postdoctoral fellowship and T.I.E. and K.A.H. acknowledges the NSF for research grants CHE-0089172 and OCE-9708478.

## ■ REFERENCES

- (1) Orr, W. L.; Grady, J. R. Perylene in Basin Sediments off Southern California. *Geochim. Cosmochim. Acta* **1967**, *31*, 1201–1209. [https://doi.org/10.1016/S0016-7037\(67\)80058-9](https://doi.org/10.1016/S0016-7037(67)80058-9).
- (2) Aizenshtat, Z. Perylene and Its Geochemical Significance. *Geochim. Cosmochim. Acta* **1973**, *37*, 559–567. [https://doi.org/10.1016/0016-7037\(73\)90218-4](https://doi.org/10.1016/0016-7037(73)90218-4).
- (3) Silliman, J. E.; Meyers, P. A.; Ostrom, P. H.; Ostrom, N. E.; Eadie, B. J. Insights into the Origin of Perylene from Isotopic Analyses of Sediments from Saanich Inlet. *Org. Geochem.* **2000**, *31* (11), 1133–1142. [https://doi.org/10.1016/S0146-6380\(00\)00120-0](https://doi.org/10.1016/S0146-6380(00)00120-0).
- (4) Zhang, X.; Xu, Y.; Ruan, J.; Ding, S.; Huang, X. Origin, Distribution and Environmental Significance of Perylene in Okinawa

Trough since Last Glaciation Maximum. *Org. Geochem.* **2014**, *46*, 288–294. <https://doi.org/10.1016/j.orggeochem.2014.09.008>.

(5) Venkatesan, M. I. Occurrence and Possible Sources of Perylene in Marine Sediments—a Review \*. *Mar. Chem.* **1988**, *25* (2845), 1–27. [https://doi.org/10.1016/0304-4203\(88\)90011-4](https://doi.org/10.1016/0304-4203(88)90011-4).

(6) Meyers, P. A.; Ishiwatari, R. Lacustrine Organic Geochemistry—an Overview of Indicators of Organic Matter Sources and Diagenesis in Lake Sediments. *Org. Geochem.* **1993**, *20* (7), 867–900. [https://doi.org/10.1016/0146-6380\(93\)90100-P](https://doi.org/10.1016/0146-6380(93)90100-P).

(7) Itoh, N.; Tamamura, S.; Kumagai, M. Distributions of Polycyclic Aromatic Hydrocarbons in a Sediment Core from the North Basin of Lake Biwa. *Org. Geochem.* **2010**, *41* (8), 845–852. <https://doi.org/10.1016/j.orggeochem.2010.04.002>.

(8) Gschwend, P. M.; Hites, R. A. Fluxes of Polycyclic Aromatic Hydrocarbons to Marine and Lacustrine Sediments in the Northeastern United States. *Geochim. Cosmochim. Acta* **1981**, *45* (12), 2359–2367. [https://doi.org/10.1016/0016-7037\(81\)90089-2](https://doi.org/10.1016/0016-7037(81)90089-2).

(9) Wilcke, W.; Krauss, M.; Amelung, W. Carbon Isotope Signature of Polycyclic Aromatic Hydrocarbons (PAHs): Evidence for Different Sources in Tropical and Temperate Environments? *Environ. Sci. Technol.* **2002**, *36* (16), 3530–3535. <https://doi.org/10.1021/es020032h>.

(10) Gocht, T.; Barth, J. A. C.; Epp, M.; Jochmann, M.; Blessing, M.; Schmidt, T. C.; Grathwohl, P. Indications for Pedogenic Formation of Perylene in a Terrestrial Soil Profile: Depth Distribution and First Results from Stable Carbon Isotope Ratios. *Appl. Geochem.* **2007**, *22* (12), 2652–2663. <https://doi.org/10.1016/j.apgeochem.2007.06.004>.

(11) Schnurmann, R.; Maddams, W. F.; Barlow, M. C. Spectrophotometric Identification of Polynuclear Aromatic Components In High Boiling Petroleum Fractions. *Anal. Chem.* **1953**, *25* (7), 1010–1013. <https://doi.org/10.1021/ac60079a003>.

(12) Wise, S. A.; Poster, D. L.; Leigh, S. D.; Rimmer, C. A.; Mössner, S.; Schubert, P.; Sander, L. C.; Schantz, M. M. Polycyclic Aromatic Hydrocarbons (PAHs) in a Coal Tar Standard Reference Material-SRM 1597a Updated. *Anal. Bioanal. Chem.* **2010**, *398* (2), 717–728. <https://doi.org/10.1007/s00216-010-4008-x>.

(13) Simoneit, B. R. T.; Rogge, W. F.; Lang, Q.; Jaffé, R. Molecular Characterization of Smoke from Campfire Burning of Pine Wood (*Pinus Elliottii*). *Chemosph. - Glob. Chang. Chemosphere: Global Change Sci.* **2000**, *2* (1), 107–122. [https://doi.org/10.1016/S1465-9972\(99\)00048-3](https://doi.org/10.1016/S1465-9972(99)00048-3).

(14) Wang, Z.; Fingas, M.; Shu, Y. Y.; Sigouin, L.; Landriault, M.; Lambert, P.; Turpin, R.; Campagna, P.; Mullin, J. Quantitative Characterization of PAHs in Burn Residue and Soot Samples and Differentiation of Pyrogenic PAH1 from Petrogenic PAHs - The 1994 Mobile Burn Study. *Environ. Sci. Technol.* **1999**, *33* (18), 3100–3109. <https://doi.org/10.1021/es990031y>.

(15) Lima, A. L.; Eglinton, T. I.; Reddy, C. M. High-Resolution Record of Pyrogenic Polycyclic Aromatic Hydrocarbon Deposition during the 20th Century. *Environ. Sci. Technol.* **2003**, *37* (1), 53–61. <https://doi.org/10.1021/es025895p>.

(16) Silliman, J. E.; Meyers, P. A.; Eadie, B. J.; Val Klump, J. A Hypothesis for the Origin of Perylene Based on Its Low Abundance in Sediments of Green Bay. *Chem. Geol.* **2001**, *177* (3–4), 309–322. [https://doi.org/10.1016/S0009-2541\(00\)00415-0](https://doi.org/10.1016/S0009-2541(00)00415-0).

(17) Wakeham, S. G.; Canuel, E. A. Biogenic Polycyclic Aromatic Hydrocarbons in Sediments of the San Joaquin River in California (USA), and Current Paradigms on Their Formation. *Environ. Sci. Pollut. Res.* **2016**, *23* (11), 10426–10442. <https://doi.org/10.1007/s11356-015-5402-x>.

(18) Fan, C. W.; Shiue, J.; Wu, C. Y.; Wu, C. Y. Perylene Dominance in Sediments from a Subtropical High Mountain Lake. *Org. Geochem.* **2011**, *42* (1), 116–119. <https://doi.org/10.1016/j.orggeochem.2010.10.008>.

(19) Hites, R. A.; Laflamme, R. E.; Windsor, J. G.; Farrington, J. W.; Deuser, W. G. Polycyclic Aromatic Hydrocarbons in an Anoxic Sediment Core from the Pettaquamscutt River (Rhode Island, U.S.A.).

*Geochim. Cosmochim. Acta* **1980**, *44* (6), 873–878. [https://doi.org/10.1016/0016-7037\(80\)90267-7](https://doi.org/10.1016/0016-7037(80)90267-7).

(20) Gschwend, P. M.; Chen, P. H.; Hites, R. A. On the Formation of Perylene in Recent Sediments: Kinetic Models. *Geochim. Cosmochim. Acta* **1983**, *47* (12), 2115–2119. [https://doi.org/10.1016/0016-7037\(83\)90036-4](https://doi.org/10.1016/0016-7037(83)90036-4).

(21) Wakeham, S. G. Synchronous Fluorescence Spectroscopy and Its Application to Indigenous and Petroleum-Derived Hydrocarbons in Lacustrine Sediments. *Environ. Sci. Technol.* **1977**, *11* (3), 272–276. <https://doi.org/10.1021/es60126a012>.

(22) Grice, K.; Lu, H.; Atahan, P.; Asif, M.; Hallmann, C.; Greenwood, P.; Maslen, E.; Tulipani, S.; Williford, K.; Dodson, J. New Insights into the Origin of Perylene in Geological Samples. *Geochim. Cosmochim. Acta* **2009**, *73* (21), 6531–6543. <https://doi.org/10.1016/j.gca.2009.07.029>.

(23) Itoh, N.; Sakagami, N.; Torimura, M.; Watanabe, M. Perylene in Lake Biwa Sediments Originating from *Cenococcum Geophilum* in Its Catchment Area. *Geochim. Cosmochim. Acta* **2012**, *95*, 241–251. <https://doi.org/10.1016/j.gca.2012.07.037>.

(24) Louda, J. W.; Baker, E. W. Perylene Occurrence, Alkylation and Possible Sources in Deep-Ocean Sediments. *Geochim. Cosmochim. Acta* **1984**, *48* (5), 1043–1058. [https://doi.org/10.1016/0016-7037\(84\)90195-9](https://doi.org/10.1016/0016-7037(84)90195-9).

(25) Jautzy, J. J.; Ahad, J. M. E.; Hall, R. I.; Wiklund, J. A.; Wolfe, B. B.; Gobeil, C.; Savard, M. M. Source Apportionment of Background PAHs in the Peace-Athabasca Delta (Alberta, Canada) Using Molecular Level Radiocarbon Analysis. *Environ. Sci. Technol.* **2015**, *49* (15), 9056–9063. <https://doi.org/10.1021/acs.est.5b01490>.

(26) Obase, K.; Douhan, G. W.; Matsuda, Y.; Smith, M. E. Progress and challenges in understanding the biology, diversity, and biogeography of *Cenococcum gophilum*. In *Biogeography of Mycorrhizal Symbiosis*; Tedersoo, L., Ed.; Springer International Publishing, 2017; Vol. *Ecological*, pp 299–318. <https://doi.org/10.1007/978-3-319-56363-3>.

(27) Suzuki, N.; Yessalina, S.; Kikuchi, T. Probable Fungal Origin of Perylene in Late Cretaceous to Paleogene Terrestrial Sedimentary Rocks of Northeastern Japan as Indicated from Stable Carbon Isotopes. *Org. Geochem.* **2010**, *41* (3), 234–241. <https://doi.org/10.1016/j.orggeochem.2009.11.010>.

(28) Strullu-Derrien, C.; Selosse, M. A.; Kenrick, P.; Martin, F. M. The Origin and Evolution of Mycorrhizal Symbioses: From Palaeomycology to Phylogenomics. *New Phytol.* **2018**, *220* (4), 1012–1030. <https://doi.org/10.1111/nph.15076>.

(29) Allport, D. C.; Bu'Lock, J. D. Biosynthetic Pathways in *Daldinia Concentrica*. *J. Chem. Soc.* **1960**, 654–662. <https://doi.org/10.1039/JR9600000654>.

(30) Wu, H.; Lao, X. F.; Wang, Q. W.; Ren-Rong, L.; Shen, C.; Zhang, F.; Liu, M.; Jia, L. The Shiraiachromes: Novel Fungal Perylenequinone Pigments from *Shiraia bambusicola*. *J. Nat. Prod.* **1989**, *52* (5), 948–951. <https://doi.org/10.1021/np50065a006>.

(31) Daub, M. E.; Herrero, S.; Chung, K. R. Photoactivated Perylenequinone Toxins in Fungal Pathogenesis of Plants. *FEMS Microbiol. Lett.* **2005**, *252* (2), 197–206. <https://doi.org/10.1016/j.femsle.2005.08.033>.

(32) Marynowski, L.; Smolarek, J.; Bechtel, A.; Philippe, M.; Kurkiewicz, S.; Simoneit, B. R. T. Perylene as an Indicator of Conifer Fossil Wood Degradation by Wood-Degrading Fungi. *Org. Geochem.* **2013**, *59*, 143–151. <https://doi.org/10.1016/j.orggeochem.2013.04.006>.

(33) Kaden, D. A.; Hites, R. A.; Thilly, W. G. Mutagenicity of soot and associated polycyclic aromatic hydrocarbons to *Salmonella tphimurium*. *Cancer Res.* **1979**, *39* (10), 4152–4159. <https://doi.org/10.1109/ICDM.2018.00085>.

(34) United States Environmental Protection Agency. *Provisional Peer Reviewed Toxicity Values for Perylene (CASRN 198–55–0)*; Cincinnati, **2006**.

(35) Hanke, U. M.; Reddy, C. M.; Braun, A. L. L.; Coppola, A. I.; Haghypour, N.; McIntyre, C. P.; Wacker, L.; Xu, L.; McNichol, A. P.; Abiven, S.; Schmidt, M. W. I.; Eglinton, T. I. What on Earth Have We

Been Burning? Deciphering Sedimentary Records of Pyrogenic Carbon. *Environ. Sci. Technol.* **2017**, *51* (21), 12972–12980. <https://doi.org/10.1021/acs.est.7b03243>.

(36) Hubeny, J. B.; King, J. W.; Cantwell, M. Anthropogenic Influences on Estuarine Sedimentation and Ecology: Examples from the Varved Sediments of the Pettaquamscutt River Estuary, Rhode Island. *J. Paleolimnol.* **2009**, *41*, 297–314. <https://doi.org/10.1007/s10933-008-9226-2>.

(37) Hanke, U. M.; Eglinton, T. I.; Braun, A. L. L.; Reddy, C. M.; Wiedemeier, D. B.; Schmidt, M. W. I. Decoupled Sedimentary Records of Combustion: Causes and Implications. *Geophys. Res. Lett.* **2016**, *43* (10), 5098–5108. <https://doi.org/10.1002/2016GL069253>.

(38) Fitzgerald, W. F.; Engstrom, D. R.; Hammerschmidt, C. R.; Lamborg, C. H.; Balcom, P. H.; Lima-Braun, A. L.; Bothner, M. H.; Reddy, C. M. Global and Local Sources of Mercury Deposition in Coastal New England Reconstructed from a Multiproxy, High-Resolution. *Environ. Sci. Technol.* **2018**, *52* (14), 7614–7620. <https://doi.org/10.1021/acs.est.7b06122>.

(39) Lima, A. L.; Hubeny, J. B.; Reddy, C. M.; King, J. W.; Hughen, K. A.; Eglinton, T. I. High-Resolution Historical Records from Pettaquamscutt River Basin Sediments: 1. 210Pb and Varve Chronologies Validate Record of 137Cs Released by the Chernobyl Accident. *Geochim. Cosmochim. Acta* **2005**, *69* (7), 1803–1812. <https://doi.org/10.1016/j.gca.2004.10.009>.

(40) Drenzek, N. J.; Montluçon, D. B.; Yunker, M. B.; Macdonald, R. W.; Eglinton, T. I. Constraints on the Origin of Sedimentary Organic Carbon in the Beaufort Sea from Coupled Molecular 13C and 14C Measurements. *Mar. Chem.* **2007**, *103* (1–2), 146–162. <https://doi.org/10.1016/j.marchem.2006.06.017>.

(41) Eglinton, T. I.; Aluwihare, L. I.; Bauer, J. E.; Druffel, E. R. M.; McNichol, A. P. Gas Chromatographic Isolation of Individual Compounds from Complex Matrices for Radiocarbon Dating. *Anal. Chem.* **1996**, *68*, 904–912. <https://doi.org/10.1021/ac9508513>.

(42) Reimer, P. J.; Brown, T. A.; Reimer, R. W. Discussion: Reporting and Calibration of Post-Bomb 14C Data. *Radiocarbon* **2004**, *46* (3), 1299–1304. [https://doi.org/10.2458/azu\\_js\\_rc.46.4183](https://doi.org/10.2458/azu_js_rc.46.4183).

(43) Reimer, P. J.; Bard, E.; Bayliss, A.; Beck, J. W.; Blackswell, P. G.; Ramsey, C. B.; Buck, C. E.; Cheng, H.; Edwards, R. L.; Friedrich, M.; Grootes, P. M.; Guilderson, T. P.; Hafliadason, H.; Hajdas, I.; Hatté, C.; Heaton, T. J.; Hoffmann, D. L.; Hogg, A. G.; Hughen, K. A.; Kaiser, K. F.; Kromer, B.; Manning, S. W.; Niu, M.; Reimer, R. W.; Richards, D. A.; Scott, E. M.; Southon, J. R.; Staff, R. A.; Turney, C. S. M.; van der Plicht, J. Intcal13 and Marine13 Radiocarbon Age Calibration Curves 0–50,000 Years Cal BP. *Radiocarbon* **2013**, *55* (4), 1869–1887. [https://doi.org/10.2458/azu\\_js\\_rc.55.16947](https://doi.org/10.2458/azu_js_rc.55.16947).

(44) Blattmann, T. M.; Letsch, D.; Eglinton, T. I. On the Geological and Scientific Legacy of Petrogenic Organic Carbon. *Am. J. Sci.* **2018**, *318* (8), 861–881. <https://doi.org/10.2475/08.2018.02>.

(45) Gaines, A. G. Papers on the Geomorphology, Hydrography and Geochemistry of the Pettaquamscutt River Estuary. PhD Dissertation, University of Rhode Island, 1975.

(46) Meyers, P. A. Preservation of Elemental and Isotopic Source Identification of Sedimentary Organic Matter. *Chem. Geol.* **1994**, *114* (3–4), 289–302. [https://doi.org/10.1016/0009-2541\(94\)90059-0](https://doi.org/10.1016/0009-2541(94)90059-0).

(47) Levin, I.; Kromer, B.; Hammer, S. Atmospheric D14CO2 Trend in Western European Background Air from 2000 to 2012. *Tellus, Ser. B* **2013**, *65* (20092), 1–7. <https://doi.org/10.3402/tellusb.v65i0.20092>.

(48) Meyers, P. A. Organic Geochemical Proxies of Paleoenvironmental, Paleolimnologic, and Paleoclimatic Processes. *Org. Geochem.* **1997**, *27* (5/6), 213–250. [https://doi.org/10.1016/S0146-6380\(97\)00049-1](https://doi.org/10.1016/S0146-6380(97)00049-1).

(49) Tan, Y. L.; Heit, M. Biogenic and Abiogenic Polynuclear Aromatic Hydrocarbons in Sediments from Two Remote Adirondack Lakes. *Geochim. Cosmochim. Acta* **1981**, *45* (11), 2267–2279. [https://doi.org/10.1016/0016-7037\(81\)90076-4](https://doi.org/10.1016/0016-7037(81)90076-4).

(50) Slater, G. F.; Benson, A. A.; Marvin, C.; Muir, D. PAH Fluxes to Siskiwit Revisited: Trends in Fluxes and Sources of Pyrogenic PAH and Perylene Constrained via Radiocarbon Analysis. *Environ. Sci. Technol.* **2013**, *47* (10), 5066–5073. <https://doi.org/10.1021/es400272z>.

(51) O’Leary, M. Carbon Isotopes in Photosynthesis. *BioScience* **1988**, *38* (5), 328–336. <https://doi.org/10.2307/1310735>.

(52) Lehmann, J.; Kleber, M. The Contentious Nature of Soil Organic Matter. *Nature* **2015**, *528*, 60–68. <https://doi.org/10.1038/nature16069>.

(53) Berhe, A. A.; Barnes, R. T.; Six, J.; Marin-Spiotta, E. Role of Soil Erosion in Biogeochemical Cycling of Essential Elements: Carbon, Nitrogen, and Phosphorus. *Annu. Rev. Earth Planet. Sci.* **2018**, *46* (1), 521–548. <https://doi.org/10.1146/annurev-earth-082517-010018>.

(54) Douglas, P. M. J.; Pagani, M.; Eglinton, T. I.; Brenner, M.; Hodell, D. A.; Curtis, J. H.; Ma, K. F.; Breckenridge, A. Pre-Aged Plant Waxes in Tropical Lake Sediments and Their Influence on the Chronology of Molecular Paleoclimate Proxy Records. *Geochim. Cosmochim. Acta* **2014**, *141*, 346–364. <https://doi.org/10.1016/j.gca.2014.06.030>.

(55) French, K. L.; Hein, C. J.; Haghipour, N.; Wacker, L.; Kudrass, H. R.; Eglinton, T. I.; Galy, V. Millennial Soil Retention of Terrestrial Organic Matter Deposited in the Bengal Fan. *Sci. Rep.* **2018**, *8* (1), 1–8. <https://doi.org/10.1038/s41598-018-30091-8>.

(56) Vonk, J. E.; Drenzek, N. J.; Hughen, K. A.; Stanley, R. H. R.; McIntyre, C.; Montluçon, D. B.; Giosan, L.; Southon, J. R.; Santos, G. M.; Druffel, E. R. M.; Andersson, A. A.; Sköld, M.; Eglinton, T. I. Temporal Deconvolution of Vascular Plant-Derived Fatty Acids Exported from Terrestrial Watersheds. *Geochim. Cosmochim. Acta* **2019**, *244*, 502–521. <https://doi.org/10.1016/j.gca.2018.09.034>.

(57) Coppola, A. I.; Wiedemeier, D. B.; Galy, V.; Haghipour, N.; Hanke, U. M.; Nascimento, G. S.; Usman, M.; Blattmann, T. M.; Reisser, M.; Freymond, C. V.; Zhao, M.; Voss, B.; Wacker, L.; Schefuß, E.; Peucker-Ehrenbrink, B.; Abiven, S.; Schmidt, M. W. I.; Eglinton, T. I. Global-Scale Evidence for the Refractory Nature of Riverine Black Carbon. *Nat. Geosci.* **2018**, *11* (8), 584–588. <https://doi.org/10.1038/s41561-018-0159-8>.

(58) Burdige, D. J. Burial of Terrestrial Organic Matter in Marine Sediments: A Re-Assessment. *Global Biogeochem. Cycles* **2005**, *19* (4), 1–7. <https://doi.org/10.1029/2004gb002368>.

(59) Oros, D. R.; Simoneit, B. R. T. *Identification and Emission Factors of Molecular Tracers in Organic Aerosols from Biomass Burning Part 1. Temperate Climate Conifers*; 2001; Vol. 16. [https://doi.org/10.1016/S0883-2927\(01\)00021-X](https://doi.org/10.1016/S0883-2927(01)00021-X).

(60) Venkatesan, M. I.; Kaplan, I. R. The Lipid Geochemistry of Antarctic Marine Sediments: Bransfield Strait. *Mar. Chem.* **1987**, *21* (2844), 347–375. [https://doi.org/10.1016/0304-4203\(87\)90056-9](https://doi.org/10.1016/0304-4203(87)90056-9).

(61) Jiang, C.; Alexander, R.; Kagi, R. I.; Murray, A. P. Origin of Perylene in Ancient Sediments and Its Geological Significance. *Org. Geochem.* **2000**, *31* (12), 1545–1559. [https://doi.org/10.1016/S0146-6380\(00\)00074-7](https://doi.org/10.1016/S0146-6380(00)00074-7).

(62) Sánchez-Ramírez, S.; Wilson, A. W.; Ryberg, M. Overview of Phylogenetic Approaches to Mycorrhizal Biogeography, Diversity and Evolution. In *Biogeography of Mycorrhizal Symbiosis*; Tedersoo, L., Ed.; Springer International Publishing, 2017; Vol. 230, pp 1–38. <https://doi.org/10.1007/978-3-319-56363-3>.

(63) Blumer, M. Organic Pigments: Their Long-Term Fate. *Science* **1965**, *149* (3685), 722–726. <https://doi.org/10.1126/science.149.3685.722>.

(64) Fernandez, C. W.; Langley, J. A.; Chapman, S.; McCormack, M. L.; Koide, R. T. The Decomposition of Ectomycorrhizal Fungal Necromass. *Soil Biol. Biochem.* **2016**, *93*, 38–49. <https://doi.org/10.1016/j.soilbio.2015.10.017>.

(65) Bianchi, T. S.; Cui, X.; Blair, N. E.; Burdige, D. J.; Eglinton, T. I.; Galy, V. Centers of Organic Carbon Burial and Oxidation at the Land-Ocean Interface. *Org. Geochem.* **2018**, *115*, 138–155. <https://doi.org/10.1016/j.orggeochem.2017.09.008>.

(66) Hedges, J. I.; Oades, J. M. Comparative Organic Geochemistries of Soils and Marine Sediments. *Org. Geochem.* **1997**, *27* (7–8), 319–361. [https://doi.org/10.1016/s0146-6380\(97\)00056-9](https://doi.org/10.1016/s0146-6380(97)00056-9).